
Reiseeindrücke

aus Sri Lanka

von

Herbert Becker



- circa 240 km östlich der Südspitze
- kürzeste Entfernung circa 55 Km
- 21 Millionen Einwohner
- 65000km3 etwas kleiner als Bayern
- bis 1972 Ceylon


Sri Lanka ist bekannt für:

seinen Tee (Ceylon – Tee)
die Produktion und den Export von Kaffee, Kautschuk und Kokosnüssen
seine Edelsteine (Ceylon – Saphir)
Die Insel ist aufgrund ihrer landschaftlichen Schönheit und ihres reichen Kulturerbes ein beliebtes Touristenziel




Club Palm Bay Hotel
45,5 km Fahrzeit 1 Sunde 15 Minuten




Kudamaduwella
Matta Kotuwa
Puttalam
B419
Kudawewa Railway
Mahawewa
Mahawawa
WEEHENA
Chilaw - Colombo Main Rd
Aanankalliya
B419
Walahapitiya Railway
Beach Rd
HALPANWILA
Walahapitiya Rd
Club Palm Bay
Church Rd
Beach Rd
HATHTHINIYA









CLUB PALM BAY
IN






















Palmenhörnchen (*Funambulus*)






Glanzkrähen (Corvus splendens)










DESKTOP
LAPTOP
TABLET
ALARM


75-2208


හලාවත
சிலாபம்
20 km
A 3
CHILAW
නාත්තන්ඩිය
நாத்தாண்டிய
4 km
B 272
NATTANDIYA
RDA











032-2255944 / 072-6530575








Club Palm Bay Hotel
2 Std. 43 Min.
124 km
2 Std. 38 Min.
120 km
Chilaw
Bingiriya
Wariyapola
Katupotha
Gokarella
Ridigama
Panagamuwa
Kuliyapitiya
Kurunegala
Matale
Narammala
Pothuhera
Wennappuwa
Dankotuwa
Rambukkana
Katugastota
Kandy
Digana
Kegalle
Mawanella
Warakapola
Negombo
Katunayake
Veyangoda
Gampola
Gampaha
Dedugala
Ruwanwella
Nawalapitiya
Ramboda
Wattala
Ginigathhena
Avissawella
Colombo
Google
Kartendaten © 2015 Google
Bedingungen
Datenschutz
Feedback geben
10 km

Die Schwarzbauch Erdschildkröte ist eine mittelgroße Süßwasserschildkröte mit zur Zeit 6 anerkannten Unterarten, von denen zwei auf Sri Lanka vorkommen

Melanochelys trijuga trijuga (Indien)
Melanochelys trijuga coronata (Indien)
Melanochelys trijuga edeniana (Myanmar)
Melanochelys trijuga indopeninsularis (Bangladesh, Indin, Nepal)
Melanochelys trijuga parkeri (Sri Lanka)
Melanochelys trijuga thermalis (Indien, Maldiven, Sri Lanka)





Hutaffen (Macaca sinica sinica)






SAVE
CROCS




































Sägerückenagamen (*Calotes calotes*)
























Kandy



B519
B489
B69
Natha Devalaya
Sri Dalada Maligawa
International Buddhist Museum
Lewella Junction
Lewella Junction
Maligawa
National Museum
Buwalikada Shed
A26
A26
Kandy City Centre
A1
Maligawa
B521
Hotel Thilanka
Lake Round
A26
Buwalikada
B551
Wales Park
Hillwood
Mahamaya Primary
Malwathu Maha Viharaya
Hotel Suisse
B195
Dharmaraja Mawatha
B551
Reservoir Rd
Rajapihilla Mawatha
Mahamaya
Lakeside
Mahamaya Girls College
Rathubokkuwa
Google





Verbreitung des Bindenwarans (Koch et al 2007, Koch & Böhme 2010)
Gelb: V. s. salvator, Grün: V. s. macromaculatus, Blau: V. s. bivittatus, Rot: V. s. andamanensis, Braun: V. s. ziegleri, Grau: noch nicht als Unterart abgegrenzte Populationen auf Sulawesi














Kormoran (*Phalacrocorax carbo sinensis*)



Riesenflughund - Kolonie (*Pteropus giganteus giganteus*)



























Udawatta Kele Sanctuary



Udawatta Kele Sanctuary

ein historisches Waldreservat auf einem Bergrücken in der Stadt Kandy. Es enthält eine große Vielfalt an Pflanzenarten, vor allem Lianen, Sträucher und kleine Bäume. Udawatta Kele wurde als Waldschutzgebiet im Jahr 1856 bestimmt.














Schwarznarbenkröte (*Duttaphrynus melanostictus*)



Pinnawala Elephantenwaisenhaus












A/L PHYSICS
අලි නාවන ස්ථානය
யானைகள் குளிக்கும் இடம்
ELEPHANT BATHING PLACE
10.00-
12.00
02.00-
04.00
POO PAPER SOLD HERE !
POO PAPER SOLD HERE !
POO PAPER SOLD HERE !










Blutsaugeragame (Calotes versicolor)

























සේවය සලසන දුම්රිය සුරකිමු.
KADUGANNAWA
KANDY
HAT TON
COLOMBO
ANURADHAPURA
MATARA
1
සාදර අනුග්‍රහය
රඹුක්කන ගෝල්ඩ් හවුස්
RGH
RAMBUKKANA GOLD HOUSE
91/3, කුරුණෑගල පාර,
රඹුක්කන.
91/3,KURUNEGALA ROAD,
RAMBUKKANA.
2


OFV - 8355


FOR CLERGY


KA 1011
airtel
CALTEX

Die größten Teeproduzenten weltweit in Tonnen (2012)

1	Volksrepublik China	1.714.902
2	Indien	1.000.000
3	Kenia	369.400
4	Sri Lanka	330.000
5	Türkei	225.000
6	Vietnam	216.900
7	Iran	158.000
8	Indonesien	150.100
9	Argentinien	100.000
10	Japan Japan	85.900


Kadugannawa Tea
& Promotion Centre




ROLL NO. 03
Double Action
MARSHALL
CEYLON


SIROCCO
DAVIDSON & Co LTD
BELFAST, IRELAND
DAVIDSON'S
PATENT
SIROCCO
AIR-HEATER




2387


2396

Royal Botanic Gardens













































hinduistischen Muthumariamman
Thevasthanam Tempel






PEPSI
PEPSI
PEPSI



ROOMS
Sarilco
NEW
SHAN
INTERNA
066
HOTEL

goldener Buddha in Dambulla








Meegalewa
Galnewa
Kekirawa
Habarana
Hingurakgoda
Avukana
Girithale
Galgamuwa
Sigiriya
Palugasdama
Polonnaruwa
Ambanpola
Inamaluwa
Dambulla
Saliyagama
Madagalla
Polpithigama
Bulanawewa
Maho
Bakamuna
Wasgamuwa
National Park
Galewela
Elahara
Padeniya
Melsiripura
Wariyapola
Gokarella
Dehiattaka
Sandunp
Nawa
Madagam
Ridigama
Laggala
Hettipola
Diyawiddag
Mahaweli Ganga
Panagamuwa
Kurunegala
Matale
Knuckles
Forest Reserve
Girandurukotte
Pothuhera
Hasalaka
Katugastota
Mahiyanganaya
Rambukkana
Kandy
Digana
Mediya
Google
A6
A9
A6
A1
A28
B89
A28
AB44
A9
A10
B409
B409
A6
AB26
A10
B15
A9
AB44
B8
A10
A26
B8
B5
A6
A19





Bronzeskink (Eutropis macularia)

































Sägerückenagame (*Calotes calotes*)









Bronzeskink (Eutropis macularia)

















Weißbartlanguren (Presbytis senex vetulus)


Giriulla
Katugastota
Mahiyanganaya
Dankotuwa
Rambukkana
Kandy
Victoria
Randenigala
Rantembe
Sanctuary
Gal Oya
National Park
Negombo
Bibile
Walapane
Lunugala
Ginigathhena
Badulla
Nuwara Eliya
Colombo
Siyambal
Ella
Dehiwala-Mount
Lavinia
Horton Plains
National Park
Bandarawela
Wattegama
Haputale
Buttala
Ratnapura
Balangoda
Wadduwa
Madampe
Udawalawe
National Park
Beruwala
Udawalawa
Kataragama
Sinharaja
Forest Reserve
Turtle Hatchery
Yala
Tissamaharama
Kirinda
Hikkaduwa
Hambantota
Galle
Google
Tangalle
A1
A5
A21
A7
A22
A8
A4
A2
A18
E03
E02
E01











Vor der Küste Sri Lankas kommen 5 Meeresschildkrötenarten vor und legen ihre Eier an den Stränden ab:

Suppenschildkröte (*Chelonia mydas*
Lederschildkröte (*Dermochelys coriacea*
Echte Karettschildkröte (*Eretmochelys imbricata*)
Unechte Karettschildkröte (*Caretta caretta*)
Oliv-Bastardschildkröte (*Lepidochelys olivacea*


Coastal Plants (Geen Belt) & Marine Turtles Information Centre
මුහුදු ශාක හා මුහුදු කැස්බෑ තොරතුරු මධ්‍යස්ථානය
මුහුදු කැස්බෑ පර්යේෂණ හා සංරක්ෂණ ව්‍යාපෘතිය
ගාලු පාර බෙන්තොට දකුණ
Sea Turtles Research and Concervation Project
Galle road Bentota South
Join With us To Protect Turtles World Wide.


WELCOME
SEA TURTLES
PROJECT
NO.390,BENTOTA SOUTH
SRILANKA










Green Turtle - Chelonia Mydas
Habitat :
Tropical and subtropical oceans in the open seas
Feeding : Seagrass and Algae






Suppenschildkröte (Chelonia mydas)


සුනාමි ඛේදවාචකය TSUNAMI DISASTER
Sea turtles conservation (2004-12-26)
Research Project Bentota South, Sri Lanka.






Green Turtel

කොළ කැස්බෑවා

Scientific name: Chelonia mydas



The green sea turtle, also known as the green turtle, black turtle, or Pacific green turtle, is a large sea turtle of the family Cheloniidae. It is the only species in the genus Chelonia. Wikipedia
Rank: Species Higher classification: Chelonl

Maximum Weight 350 kg , Max. Le. 1.2m

කොළ කැස්බෑවා කුඩා අවධියේ මාංශ භක්ෂකයෙකු වුවද පරිනත අවස්ථාවේදී ශාක භක්ෂකයකු වේ
බර කිලෝ ග්‍රෑම් 350 ක් පමණ වේ දිගින් මීටර 1 1/4 පමණ වේ වසර 100 කට අධික කාලයක් ජීවත් වේ
වරකට බිත්තර 150 ක් පමණ දමයි බිත්තරයක බර 38g පමණ වේ




SEA TURTLES PROJECT
BENTOTA SOUTH , SRI LANKA




SEA TURTLES CONSERVATION RESEARCH PROJECT
BENTOTA, SRI LANKA
TURTLES EGGS HATCHERY
The Green Turtles
(Chelonia Mydas)
The Olive Ridley Turtles
(Lepidochelys Olivacea)
The Hawksbill Turtles
(Eremochelys Imbricat)
The Loggerhead Turtles
(Caretta Caretta)
The Leatherback Turtles
(Dermochelys Coriacea)
Name of Turtles
Weight of Egg
Green Turtles 36g
Olive Ridley Turtles 39g
Hawksbill Turtles 28g
Loggerhead Turtles 36g
Leatherback Turtles 75g
SEA TURTLES PROJECT
BENTOTA, SRI LANKA



තක්කඩ
THAKKADA






Modell der Lederschildkröte (Dermochelys coriacea)








Kosgoda Turtle Hatchery
Leatherback Turtle
Kosgoda Turtle Hatchery
Loggerhead Turtle
Kosgoda Turtle Hatchery
Hawksbill Turtle
Kosgoda Turtle Hatchery
Green Turtle
Kosgoda Turtle Hatchery
Olive ridley Turtle


Sea Turtle Identification Key
Leatherback
Dermochelys
coriacea
Green
Chelonia mydas
Black*
Flatback
Natator
Depressus
Hawksbill
Eretmochelys
imbricata
Loggerhead
Caretta caretta
Kemp's Ridley
Lepidochelys
kempii
Olive Ridley
Lepidochelys
olivacea
Leathery, no scutes;
5 long ridges
One pair prefrontal scales
4 costal (lateral) scutes;
First costal scute does not touch nuchal
Two pairs
prefrontal scales;
overlapping
scutes
5 or 6 costal scutes;
Carapace not circular
5 or more costal (lateral) scutes;
First costal scute touches nuchal
Carapace wide and
almost circular
5 costal scutes;
Nearly circular
usu. 6 or more
costal scutes
Restricted primarily to the
eastern Pacific Ocean;
Dark pigmentation
Found in tropical
coastal waters
of Australia;
Carapace has
upturned edges
Restricted primarily to
the Gulf of Mexico and
Atlantic coast of the USA
Found in tropical waters
of the Pacific, Indian and
South Atlantic Oceans
LEO


Sea Turtle Life Cycle












2014.11.02
GREEN TURTLE
Eggs 80
GREEN TURTLE
Eggs 89
GREEN TURTLE
Eggs 73
2014.11.02
GREEN TURTLE
Eggs 100
GREEN TURTLE
Eggs 70
2014.11.06
Olive Ridley
23
Eggs 130
GREEN TURTLE
Eggs 79
2014.11.03
GREEN TURTLE
Eggs 68






TAKE ME
ON A


Suppenschildkröte (Chelonia mydas)

Oliv-Bastardschildkröte (Lepidochelys olivacea)



Echte Karettschildkröte *(Eretmochelys imbricata)*



